

5-Hydroxytryptamine (Serotonin) Receptor 6 (HTR6) ACTOne™ Stable Cell Line


CATALOG NUMBER: CL-01-HTR6

Introduction

HTR6 belongs to the seven-transmembrane G protein-coupled receptor family of proteins. The encoded protein couples with the Gs alpha subunit and stimulates adenylate cyclase to activate the cyclic AMP-dependent signaling pathway. This receptor is thought to regulate cholinergic neuronal transmission in the brain. Several antidepressants and antipsychotic drugs have a high affinity for this receptor.

Description

Human HTR6 ACTOne™ is a HEK-293 CNG cell line that expresses recombinant human HTR6. HEK-293 CNG cells express a modified CNG (Cyclic Nucleotide Gated) channel that opens in response to elevated intracellular cAMP levels and consequently result in ion flux (often detectable by calcium-responsive dye, Cat# CA-C155) and cell membrane depolarization which can be easily measured with fluorescent Membrane Potential Dye (Cat# CA-M165). The assay allows both end-point and kinetic measurement of intracellular cAMP changes with a FLIPR, or a fluorescence microplate reader.



Parental Cells

HEK-293 CNG cells (originally developed by BD Biosciences by introducing CNG in HEK-293 cells) (Cat# CL-03-PC20)

Gene/Enzyme Introduced

HTR6 (Genbank Accession No. P50406)

Applications

- cAMP dependent human HTR6 cell based assay
- cell based high-throughput screening of human HTR6 inhibitors

Functional Test

- this cell line has been tested positive for HTR6 specific response

- surviving rate: More than 2.5 million/vial on the second day after thawing
- the receptor specific activity is stable for 10 weeks continuous passage

Mycoplasma Contamination Test

This lot of cells has been tested and found to be free of mycoplasma contamination.

Content

- Stable cells: 1 mL (1 x 10⁶ cells/mL in 70% DMEM, 20% FBS, 10% DMSO)

Growth Properties

Adherent

Cell Culture Medium

- Growth medium: DMEM-10% FBS supplemented with 250 µg/ml G418, 1 µg/ml Puromycin
- Freezing medium: 10% DMSO, 90% complete cell culture medium

Storage

Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below -130°C, preferably in liquid nitrogen vapor, until ready for use.

Data Analysis




Figure 1. Response of ACTOne™ HTR6 cell line & parental cell line to 5-HT.

ACTOne™ HTR6 cells and parental cells (Cat# CL-03-PC20) were plated overnight in 20 µl culture medium on a 384 well Biocoat plate. The next day, cells were dye-loaded with 20 µl/well of 1x Dye-loading solution (membrane potential dye kit, Cat# CA-M165). After 2 hours of incubation at room temperature, two readings were obtained prior to and 30 min after the addition of 5-HT. Ratios of the two readings (F/F₀) are plotted in the figure.

- Dose response curve of 5-HT H in ACTOne™ HTR6 cell line. EC₅₀ = 1.04 nM in the presence of PDE inhibitor Ro 20-1724.**
- Parental cells do not respond to 5-HT.**

Notice to Purchaser

This cell line is sold with a use license. It may not be transferred to third parties, resold, modified for resale without written approval of eEnzyme LLC. Refer to the license agreements for details on the usage restrictions.