

## Neuropeptide Y Receptor Y4 (NPY4R) ACTOne™ Stable Cell Line


CATALOG NUMBER: CL-11-NPY4R

### Introduction

Neuropeptide Y receptor Y4 (NPY4R), also called pancreatic polypeptide receptor 1 (PPYR1), is a member of the Gi/o-protein-coupled receptor family that are currently divided into four subtypes: Y1, Y2, Y4 and Y5. NPY receptors mediate a diverse range of biological actions including stimulation of food intake, anxiolysis, modulation of circadian rhythm, pain transmission and control of pituitary hormone release. All four NPY receptor subtypes are found at high concentrations in the brain and colon and are also found in a wide variety of peripheral tissues at lower levels.

### Description

Human NPY4R ACTOne™ is a HEK-293 CNG cell line that expresses recombinant human NPY4R. HEK-293 CNG cells express a modified CNG (Cyclic Nucleotide Gated) channel that opens in response to elevated intracellular cAMP levels and consequently result in ion flux (often detectable by calcium-responsive dye, Cat# CA-C155) and cell membrane depolarization which can be easily measured with fluorescent Membrane Potential Dye (Cat# CA-M165). The assay allows both end-point and kinetic measurement of intracellular cAMP changes with a FLIPR, or a fluorescence microplate reader.



### Parental Cells

HEK-293 CNG cells (originally developed by BD Biosciences by introducing CNG in HEK-293 cells) (Cat# CL-03-PC20)

### Gene/Enzyme Introduced

NPY4R (Genbank Accession No. NP\_005963)

### Applications

- cAMP dependent assay for Gi-coupled human NPY4R
- cell based high-throughput screening of human NPY4R inhibitors

### Functional Test

- this cell line has been tested positive for NPY4R specific response



- surviving rate: More than 2.5 million/vial on the second day after thawing
- the receptor specific activity is stable for 10 weeks continuous passage

## Mycoplasma Contamination Test

This lot of cells has been tested and found to be free of mycoplasma contamination.

## Content

- Stable cells: 1 mL ( $1 \times 10^6$  cells/mL in 70% DMEM, 20% FBS, 10% DMSO)

## Growth Properties

Adherent


## Cell Culture Medium

- Growth medium: DMEM-10% FBS supplemented with 250  $\mu$ g/ml G418, 1  $\mu$ g/ml Puromycin
- Freezing medium: 10% DMSO, 90% complete cell culture medium

## Storage

Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below  $-130^{\circ}\text{C}$ , preferably in liquid nitrogen vapor, until ready for use.

## Data Analysis



**Figure 1. Response of ACTOne™ NPY4R cell line & parental cell line to hPP.**

ACTOne™ NPY4R cells and parental cells (Cat# CL-03-PC20) were plated overnight in 20  $\mu$ l culture medium on a 384 well Biocoat plate. The next day, cells were dye-loaded with 20  $\mu$ l/well of 1x Dye-loading solution (membrane potential dye kit, Cat# CA-M165). After 2 hours of incubation at room temperature, two readings were obtained prior to and 30 min after the addition of hPP. Ratios of the two readings (F/F0) are plotted in the figure.

- Dose response curve of hPP in ACTOne™ NPY4R cell line. EC50 = 816 pM in the presence of PDE inhibitor Ro 20-1724 and  $\beta$ -adrenoceptor agonist isoproterenol.**
- Parental cells do not respond to hPP.**

## Notice to Purchaser

This cell line is sold with a use license. It may not be transferred to third parties, resold, modified for resale without written approval of eEnzyme LLC. Refer to the license agreements for details on the usage restrictions.

