

ACTOne™ Human Sphingosine-1-Phosphate Receptor 2 (S1PR2) Stable Cell Line

Catalog Number: CL-11-S1PR2

Product Type: Stable Cell Line

Cell Line Name: ACTOne™ S1PR2

Host Cell Line: HEK-293 CNG


Target Receptor: Human Sphingosine-1-Phosphate Receptor 2 (S1PR2)

GenBank Accession: NP_001391.2

Product Description

The ACTOne™ S1PR2 stable cell line is a HEK-293 CNG derivative engineered to express recombinant human sphingosine-1-phosphate receptor 2 (S1PR2). These cells contain a modified cyclic nucleotide-gated (CNG) channel that opens in response to intracellular cAMP elevation, triggering ion flux and membrane depolarization. This response is quantifiable using fluorescent membrane potential dye (Cat# CA-M165) or calcium-responsive dye (Cat# CA-C155).

The assay supports both endpoint and kinetic measurements of cAMP signaling and is compatible with FLIPR and fluorescence microplate readers



Applications

- Functional cAMP assays for Gi-coupled S1PR2
- High-throughput screening of S1PR2 agonists and antagonists
- GPCR pathway analysis

Functional Validation

- **S1PR2-specific response:** Confirmed
- **Post-thaw viability:** >2.5 million cells/vial on Day 2
- **Receptor activity stability:** Maintained over 10 weeks of continuous passage
- **Mycoplasma status:** Negative (tested lot-specific)

Product Format

- **Volume:** 1 mL
- **Cell Density:** 1×10^6 cells/mL
- **Cryopreservation Medium:** 70% DMEM, 20% FBS, 10% DMSO

Growth Characteristics

- **Culture Type:** Adherent
- **Growth Medium:** DMEM + 10% FBS, supplemented with 250 μ g/mL G418 and 1 μ g/mL Puromycin
- **Freezing Medium:** 10% DMSO in complete growth medium

Subculturing Protocol

1. Thaw Cryovial in 37 °C water bath (1-2 min).
2. Decontaminate vial with 70% ethanol.
3. Transfer contents to a 75 cm² flask with 20 ml of complete DMEM growth medium.
4. Replace medium after 24 hours.
5. Remove and discard culture medium next day and then add fresh DMEM complete medium.
6. Monitor daily; passage at 90% confluence (1:3 split).
7. Use 0.25% trypsin-0.53 mM EDTA for detachment (15-20 min at 37°C).
8. Centrifuge at 250 x g for 5 to 10 min.
9. Resuspend and seed at 4-6 x 10⁴ viable cells/cm².
10. Incubate at 37°C, 5% CO₂.

Freezing & Storage Instructions

1. Trypsinize cells and resuspend in freezing medium at 2.5 x 10⁶ cells/mL.
2. Aliquot 1 mL per cryovial.
3. Freeze overnight at -80°C in a cryo container.
4. Transfer to vapor-phase liquid nitrogen ($\leq -130^\circ\text{C}$) for long-term storage.



Data Summary




Figure 1. Response of ACTOne™ S1PR2 cell line & parental HEK-293 CNG cell line to S1P

ACTOne™ S1PR2 cells and parental HEK-293 CNG cells were plated overnight in 20 μ l culture medium on a 384 well Biocoat plate. The next day, cells were dye-loaded with 20 μ l/well of 1x Dye-loading solution (membrane potential dye kit, Cat# CA-M165). After 2 hours of incubation at room temperature, two readings were obtained prior to and 30 min after the addition of S1P. Ratios of the two readings (F/F₀) are plotted in the figure.

- A. Dose response curve of S1P in ACTOne™ S1PR2 cell line. EC₅₀ = 152 nM in the presence of PDE inhibitor Ro 20-1724 and β -adrenoceptor agonist isoproterenol.**
- B. Parental cells do not respond to S1P.**

Notice to Purchaser

This product is for research use only. It may not be resold, transferred, or used for commercial manufacturing or service provision without written consent from eEnzyme LLC.