

Somatostatin Receptor 5 (SSTR5) ACTOne™ Stable Cell Line


CATALOG NUMBER: CL-11-SSTR5

Introduction

SSTR5 is a multi-pass membrane protein and belongs to the G-protein coupled receptor 1 family. The activity of this receptor is mediated by G proteins which inhibit adenylyl cyclase, and different regions of this receptor molecule are required for the activation of different signaling pathways.

Description

Human SSTR5 ACTOne™ is a HEK-293 CNG cell line that expresses recombinant human SSTR5. HEK-293 CNG cells express a modified CNG (Cyclic Nucleotide Gated) channel that opens in response to elevated intracellular cAMP levels and consequently result in ion flux (often detectable by calcium-responsive dye, Cat# CA-C155) and cell membrane depolarization which can be easily measured with fluorescent Membrane Potential Dye (Cat# CA-M165). The assay allows both end-point and kinetic measurement of intracellular cAMP changes with a FLIPR, or a fluorescence microplate reader.



Parental Cells

HEK-293 CNG cells (originally developed by BD Biosciences by introducing CNG in HEK-293 cells) (Cat# CL-03-PC20)

Gene/Enzyme Introduced

SSTR5 (Genbank Accession No. NP_001044)

Applications

- cAMP dependent assay for Gi-coupled human SSTR5
- cell based high-throughput screening of human SSTR5 inhibitors

Functional Test

- this cell line has been tested positive for SSTR5 specific response
- surviving rate: More than 2.5 million/vial on the second day after thawing



- the receptor specific activity is stable for 10 weeks continuous passage

Mycoplasma Contamination Test

This lot of cells has been tested and found to be free of mycoplasma contamination.

Content

- Stable cells: 1 mL (1×10^6 cells/mL in 70% DMEM, 20% FBS, 10% DMSO)

Growth Properties

Adherent

Cell Culture Medium

- Growth medium: DMEM-10% FBS supplemented with 250 μ g/ml G418, 1 μ g/ml Puromycin
- Freezing medium: 10% DMSO, 90% complete cell culture medium

Storage

Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below -130°C , preferably in liquid nitrogen vapor, until ready for use.

Data Analysis




Figure 1. Dose response curve of Somatostatin (SST) in ACTOne™ SSTR5 cell line and parental cell line.

ACTOne™ SSTR5 cells and parental cells (Cat# CL-03-PC20) were plated overnight in 20 μ l culture medium on a 384 well Biocoat plate. The next day, cells were dye-loaded with 20 μ l/well of 1x Dye-loading solution (membrane potential dye kit, Cat# CA-M165). After 2 hours of incubation at room temperature, two readings were obtained prior to and 30 min after the addition of SST. Ratios of the two readings (F/F0) are plotted in the figure.

- Dose response curve of SST in ACTOne™ SSTR5 cell line. EC50 = 2.81 nM in the presence of PDE inhibitor Ro 20-1724 and β -adrenoceptor agonist isoproterenol.
- Parental cells do not respond to SST.

Notice to Purchaser

This cell line is sold with a use license. It may not be transferred to third parties, resold, modified for resale without written approval of eEnzyme LLC. Refer to the license agreements for details on the usage restrictions.

