

HA2 (B/Brisbane/60/2008)(aa 361-547)

CATALOG NUMBER: IB-HA2-608P, 50 µg

Introduction

Influenza hemagglutinin (HA) is a type of hemagglutinin found on the surface of the influenza viruses. HA is an antigenic glycoprotein, like all other hemagglutinins, it causes red blood cells to agglutinate. HA is responsible for binding the virus to the cell that is being infected. HA proteins bind to cells with sialic acid on the membranes, such as cells in the upper respiratory tract or erythrocytes.

HA is a homotrimeric integral membrane glycoprotein. HA monomer is synthesized as a single polypeptide that is subsequently cleaved into two smaller polypeptides, the HA1 and HA2 subunits. Each HA monomer consists of a long, helical chain anchored in the membrane by HA2 and topped by a large HA1 globule.

Description

Viral protein purified from 293 cell culture

Viral Protein

C-terminal Fc tagged HA2 (B/Brisbane/60/2008) protein (amino acid 361-547) (GenBank Accession No. ACN29380)

Applications

Western blot standard, antibody ELISA, antigen, etc.

Storage

Store at -20 °C; Stable for 6-months from the date of shipment when kept at 4 °C. Non-hazardous. No MSDS required.

Concentration


1 µg/µl in PBS

Endotoxin Level

<0.01 EU per 1 µg of the protein by LAL test

Purity

≥ 95% purity (SDS PAGE)



SDS-PAGE: purified HA2 (B/Brisbane/60/2008)(aa 361-547) protein with Fc tag

HA2 SEQ:

ERGFFGAIAGFLEGGWEGMIAGWHGYTSHGAHGVAVAADLKSTQEAINKITKNLNSLSELEVKNLQRLSGAMDELHNEILELDEKVDDLADTISQIE
LAVLLSNENGI INSEDEHLLALERKLLKMLGPSAVEI GNGCFETKHKCNQTCLDRIAAGTFDAGEFSLPTFDSLNI TAASLNDDGLDNH

hFc fragment:

SGGGSGGGSGGGSGGGSRSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNS
TYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPV
LDSGDSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKLSLSLSPGK