

Anti-Taq Monoclonal Antibody

CAT. NO. MA-029-0250 / MA-029-2500

APPLICATION

- Real-time or regular hot start PCR applications;
- PCR diagnostics, genotyping, SNP etc.

DESCRIPTION

Anti-Taq monoclonal antibodies were derived from a hybridoma (fusion of mouse myeloma cell and the cells after mouse immunization with Taq DNA Polymerase).

Anti-Taq monoclonal antibody inhibits polymerase activity before the onset of thermal cycling, preventing nonspecific amplification and primer-dimer formation. When the temperature is raised, the antibody is quickly inactivated and PCR proceeds.

Anti-Taq monoclonal antibody is effective with a variety of commercially available Taq DNA polymerases (native or recombinant). The use of hot start anti-Taq monoclonal antibody significantly improves the specificity of PCR amplification what is especially important for PCR-based diagnostics.

CLONE

8C1

CONCENTRATION

100 µg / 1 mg (4 mg/ml)

STORAGE BUFFER

10 mM Tris-HCl (pH 7.0), 50 mM KCl, 0.1 mM EDTA, 50% Glycerol

STORAGE TEMPERATURE

For long-term storage at -20 °C, stable for up to 2 years. For immediate use, keep at 4 °C, stable for up to one month.

SHIPPING CONDITIONS

Recommended shipping temperature at 4 °C or ambient condition

REACTION BUFFER

The anti-Taq monoclonal antibody reaction buffer is the same buffer used for the thermostable DNA polymerase

PURITY

> 95% by SDS-PAGE

ASSOCIATED ACTIVITIES

No conversion to the covalently closed circular DNA to the nicked or linear form was observed after incubation of 1 µg of pUC19 with antibodies in final concentration of 6 u/µl in 20 µl of reaction mixture containing 25 mM Tris-HCl (pH 7.9), 100 mM NaCl, 10 mM MgCl₂ after 16 hours at 37°C.

PROTOCOL

1. Before making any dilution of anti-Taq antibody and Taq polymerase, add 1 µg anti-Taq monoclonal antibody to 2 µg Taq DNA polymerase (~20 Units).
2. Mix gently and incubate at RT for about 10-20 minutes or store at 4°C.
3. Set up the PCR reaction by following the protocol of a regular thermal cycling condition used for the Taq DNA polymerase.

NOTE

For each Taq DNA polymerase or Taq from different batch, the antibody titration test must be performed to find the best ratio of antibody to Taq DNA polymerase.

Too much antibody added to Taq DNA polymerase could kill the PCR reaction!