

## Phosphodiesterase 2A (PDE2A) ACTOne™ Stable Cell Line


CATALOG NUMBER: CL-03-PDE2A

### Introduction

PDE2A is a cyclic nucleotide phosphodiesterase with a dual-specificity for the second messengers cAMP and cGMP, which are key regulators of many important physiological processes.

### Description

Human PDE2A ACTOne™ is a HEK293-CNG-Gs cell line that expresses recombinant human PDE2A. HEK293-CNG-Gs cells express a modified CNG (Cyclic Nucleotide Gated) channel that opens in response to elevated intracellular cAMP levels and consequently result in ion flux (often detectable by calcium-responsive dye, Cat# CA-C155) and cell membrane depolarization which can be easily measured with fluorescent Membrane Potential Dye (Cat# CA-M165). The assay allows both end-point and kinetic measurement of intracellular cAMP changes with a FLIPR, or a fluorescence microplate reader.



### Parental Cells

HEK-293 CNG-Gs cells (originally developed by BD Biosciences by introducing Gs-GPCR in HEK-293 CNG cells) (Cat# CL-03-PC10)

### Gene/Enzyme Introduced

PDE2A (Genebank Accession No. NP\_001137311.1)

### Applications

- cAMP dependent human PDE2A cell based assay

- cell based high-throughput screening of human PDE2A inhibitors

#### Functional Test

- this cell line has been tested positive for PDE2A specific response
- surviving rate: More than 2.5 million/vial on the second day after thawing
- the receptor specific activity is stable for 10 weeks continuous passage

#### Mycoplasma Contamination Test

This lot of cells has been tested and found to be free of mycoplasma contamination.

#### Content

- Stable cells: 1 mL ( $1 \times 10^6$  cells/mL in 70% DMEM, 20% FBS, 10% DMSO)
- Parental cells: 1 mL ( $1 \times 10^6$  cells/mL in 70% DMEM, 20% FBS, 10% DMSO)

#### Storage

Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below  $-130^{\circ}\text{C}$ , preferably in liquid nitrogen vapor, until ready for use.

#### Materials Required (but not supplied)

- Elite<sup>™</sup> Fluorescent Membrane Potential Dye Kit (Cat# CA-M165)
- Elite<sup>™</sup> Non-Wash Calcium Dye Assay Kit (cat# CA-C155)
- 96 or 384-well microplates: Tissue culture microplate with black wall and clear bottom is recommended.
- FlexStation (Molecular Device)

#### Cell Culture Medium

- DMEM-10% FBS supplemented with 250  $\mu\text{g/ml}$  G418, 1  $\mu\text{g/ml}$  Puromycin and 5  $\mu\text{g/ml}$  blasticidin.
- Freezing medium: 10% DMSO, 90% complete cell culture medium

#### Growth Properties


Adherent

#### Subculturing Procedure

1. Thaw the frozen cryovial of cells within 1-2 min by gentle agitation in a  $37^{\circ}\text{C}$  water bath. Decontaminate the cryovial by wiping the surface of the vial with 70% ethanol and transfer into a  $75\text{ cm}^2$  flask with 20 ml of complete DMEM growth medium.
2. Remove and discard culture medium next day, and then add fresh DMEM complete medium.
3. Monitor cell density daily. Cells should be passaged (1:3) when the culture reaches 90% confluence. Expected cell yield is between  $1.5 \times 10^5$  and  $2 \times 10^5$  viable cells/ $\text{cm}^2$ .
4. Add 2.0 to 3.0 mL of 0.25% (w/v) trypsin-0.53 mM EDTA solution to the flask and observe cells under an inverted microscope until the cell layer is dispersed (usually within 15 to 20 minutes).  
**Note:** To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Place at  $37^{\circ}\text{C}$  to facilitate dispersal.
5. Transfer cell suspension to a 15mL centrifuge tube and spin at approximately  $250 \times g$  for 5 to 10 minutes.
6. Discard supernatant and resuspend cells in fresh growth medium. Add appropriate aliquots of the cell suspension to new culture vessels. An inoculum of  $4$  to  $6 \times 10^4$  viable cells/ $\text{cm}^2$  is recommended.
7. Incubate cultures at  $37^{\circ}\text{C}$  (5%  $\text{CO}_2$ ).



Data Analysis



**Figure 1. Response of ACTOne™ PDE2A cell line & parental cell line to Bay 60-7550.**

ACTOne™ PDE2A cells and parental cells (Cat# CL-03-PDE2A) were plated overnight in 20  $\mu\text{l}$  culture medium on a 384 well Biocoat plate. The next day, cells were dye-loaded with 20  $\mu\text{l}$ /well of membrane potential dye (Cat# CA-M165). After 2 hour of incubation at room temperature, baseline was recorded using a FlexStation (Molecular Devices) (F<sub>0</sub>). 10  $\mu\text{l}$  of PDE inhibitors at various concentrations were added to the cell plate, and the data was recorded 30 minutes (F<sub>t</sub>) after drug addition. Dose response curves were generated by Prism.

**A. Dose response curve of Bay 60-7550 in ACTOne™ PDE2A cell line. EC<sub>50</sub> = 17.8 nM in the presence of 10  $\mu\text{M}$  of Ro20-1724**

**Notice to Purchaser**

This cell line is to be used for research purposes only. It may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products or to provide a service to third parties without written approval of eEnzyme LLC.